INSPECTORLAB
3301 N.W. 55TH ST., FT. LAUDERDALE, FL 33309
888-854-0477

PREPARED FOR: WILD BLUE GROUP LLC

TEST ADDRESS: 8090 RUNNING VALLEY STROUDSBURG, PA 18360-8841

CERTIFICATE OF MOLD ANALYSIS

PREPARED FOR:

WILD BLUE GROUP LLC

PHONE NUMBER: (570) 801-1300

EMAIL: THOM@WILDBLUEGROUP.COM

TEST LOCATION:

THOM SCHIAVONE

8090 RUNNING VALLEY

STROUDSBURG, PA 18360-8841

CHAIN OF CUSTODY # 52034724

COLLECTED: MON JULY 29, 2019

RECEIVED: MON JULY 29, 2019

REPORTED: MON JULY 29, 2019

APPROVED BY: JOHN D. SHANE PHD
LABORATORY MANAGER

VERSION: 1.0 (A VERSION NUMBER GREATER THAN ONE (1) INDICATES THAT THE DATA IN THIS REPORT HAS BEEN AMENDED)

EPA regulations or standards for airborne or surface mold concentrations have not been established. There are also no EPA regulations or standards for evaluating health effects due to mold exposure. Information about mold can be found at www.epa.gov/mold.

All samples were received in an acceptable condition for analysis unless noted specifically in the Comments section under a particular sample. All results relate only to the samples submitted for analysis.

A version greater than 1.0 indicates that the lab report has been revised.

IF YOU HAVE QUESTIONS REGARDING THIS REPORT, PLEASE CONTACT INSPECTORLAB AT (888) 854-0477 OR EMAIL ASK@INSPECTORLAB.COM.

INSPECTORLAB
3301 N.W. 55TH ST., FT. LAUDERDALE, FL 33309
888-854-0477

PREPARED FOR: WILD BLUE GROUP LLC **TEST ADDRESS:** 8090 RUNNING VALLEY STROUDSBURG, PA 18360-8841

Detailed Mold Report

(WATER-INDICATING FUNGI, IF PRESENT, ARE SHOWN BELOW IN RED)

Analysis Method	Air Analysis	Air Analysis	Air Analysis	Surface Analysis
Lab Sample #	52034724-1	52034724-2	52034724-3	52034724-4
Sample Identification	21023444	21045322	21042361	210423632
Sample Location	OUTSIDE CONTROL	MASTER BEDROOM	TV ROOM	TV ROOM CLOSET WALL
Sample Type / Metric	Air-O-Cell/150.0L	Air-O-Cell/150.0L	Air-O-Cell/150.0L	Swab
Analysis Date	Wed June 10, 2015	Wed June 10, 2015	Wed June 10, 2015	Wed June 10, 2015
Determination	**CONTROL**	**NORMAL**	**PROBLEM**	**GROWTH**

Fungal Types Identified	Raw Count	Spores / m^3	% of Total	Raw Count	Spores / m^3	% of Total	Raw Count	Spores / m^3	% of Total		Mold Present	
✳INDOOR PROBLEM FUNGI												
Cladosporium sphaerospermum	---	---	---	---	---	---	---	---	---		X	
Penicillium	---	---	---	---	---	---	---	---	---		X	
Penicillium/Aspergillus	---	---	---	---	---	---	1,625	10,888	66		X	
Scopulariopsis	---	---	---	---	---	---	771	5,166	31		X	
✳✳Non-Problem Fungi												
Alternaria	7	47	2	---	---	---	3	20	<1		---	
Ascospores	19	127	7	17	114	9	---	---	---		---	
Basidiospores	47	315	17	7	47	4	9	60	<1		---	
Bipolaris/Drechslera	5	34	1	1	7	<1	---	---	---		---	
Cladosporium	124	831	47	37	248	21	3	20	<1		---	
Curvularia	7	47	2	1	7	<1	---	---	---		---	
Epicoccum	6	40	2	2	13	1	---	---	---		---	
Nigrospora	4	27	1	---	---	---	---	---	---		---	
Penicillium/Aspergillus	6	40	2	91	610	52	✳	✳	✳		✳	
Pithomyces	6	40	2	2	13	1	---	---	---		---	
Smut/Myxomycetes	31	208	11	14	94	8	17	114	<1		---	
Total Spore Count	262	1,756	100	172	1,153	100	2,428	16,268	100		X	
Minimum Detection Limit		7			7			7			N/A	

Comments/Definitions	CONTROL samples are normally taken outside a building to provide a baseline from which samples on the interior of the building are compared. Outside air is considered normal whatever the mold counts may be. Light Debris: The debris present in the sample likely had no effect on the accuracy of the mold count.	Mold counts are within a NORMAL RANGE and there is no indication, based on the mold counts, that there is any exposure concern to the occupants. The LIGHT DEBRIS present in the sample likely had no effect on the accuracy of the mold count.	Mold concentrations in the air are ABNORMAL and based on the mold counts, you likely have a mold source from which spores are able to become airborne and are an exposure concern to the occupants. MODERATE DEBRIS: in the sample likely had limited affect on the accuracy of the mold count.	Presence of current or former MOLD GROWTH observed. EXPOSURE TO SPORES LIKELY and will continue if the growth is not removed. An active or intermittent water source will cause the mold to continue to grow if the water source is not eliminated.

Raw Count: Actual number of spores observed and counted.
Spores/m^3: Spores per cubic meter.
% of Total: Percentage of a particular spore in relation to total number of spores.
X: Spore type was observed.
---: Spore type was not observed.

* **Indoor Problem Fungi** are generally capable of growing on wetted building materials.

** **Non-Problem Fungi** are less capable or do not grow on wetted building materials. They are commonly found in the air outside and infiltrate into indoor air naturally. High numbers of any one of these spore types as compared to the Control sample may indicate that they are growing on wetted building materials indoors.

Spore types not listed in this report were not observed.

Background debris estimates the amount of non-spore particles. Increasing amount of debris will affect the accuracy of the spore counts. Total percent may not equal 100% due to rounding.

INSPECTORLAB
3301 N.W. 55TH ST., FT. LAUDERDALE, FL 33309
888-854-0477

Mold Glossary

PREPARED FOR: WILD BLUE GROUP LLC

TEST ADDRESS: 8090 RUNNING VALLEY STROUDSBURG, PA 18360-8841

Introduction

All spores found in indoor air are also normally found in outdoor air because most originate or live in the soil and on dead or decaying plants. Therefore, it is not unusual to find mold spores in indoor air. This Mold Glossary is only intended to provide general information about the mold found in the samples that were provided to the laboratory.

Alternaria

Outdoor Habitat: One of the most commonly observed spores in the outdoor air worldwide, normally in low numbers.

Indoor Habitat: Capable of growing on a wide variety of substrates and manufactured products found indoors when wetted.

Allergy Potential: Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis), Common cause of extrinsic asthma

Disease Potential: Not normally considered a pathogen, but can become so in immunocompromised persons.

Toxin Potential: Several known

Comments: One of the most common and potent allergens in the indoor and outdoor air. Seen in indoor air in low concentrations, probably as a result of outdoor air infiltration and/or recycling of settled dust.

Ascospores

Outdoor Habitat: Soil and decaying vegetation, dead and dying insects. These spores constitute a large part of the spores in the air and can be found in the air in very large numbers in the spring and summer, especially during and up to three (3) days after a rain.

Indoor Habitat: Very few of fungi that produce ascospores grow indoors. Some fungi that produce ascospores are recognizable by their spores and when observed are listed under their own categories. Wetted wood and gypsum wallboard paper

Allergy Potential: Depends on the type of fungus producing the ascospores.

Disease Potential: Not normally pathogenic as a group

Toxin Potential: None known

Comments: Ascospores are produced from a very large group of fungi. Notable ascospores that are considered problematic for indoor environments are Chaetomium, Peziza, and Ascotricha. If these types of ascspores are observed they will be listed in the report under their own names.

Basidiospores

Outdoor Habitat: These are mushroom spores and are common everywhere outside, especially in the late summer and fall.

Indoor Habitat: Mushrooms can grow on very wet wood products, especially on footer plates, basements, and crawlspaces. Sometimes mushrooms can be observed growing in potted plants indoors.

Allergy Potential: Rarely reported, but some Type I (hay fever, asthma) and Type III (hypersensitivity pneumonitis) has been reported.

Disease Potential: None known

Toxin Potential: None known

Comments: Mushroom spores are commonly found indoors, especially when the outdoor spore count is high. When spores of this group are derived from wood rotting fungi, including dry rot (Serpula and Poria), they can be especially destructive to buildings. When spores from destructive types of mushrooms (dry and wet rot group) are observed in the sample they are listed under their own names on the report.

Bipolaris/Drechslera

Outdoor Habitat: Commonly observed spores in the outdoor air worldwide, normally in low numbers.

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: Type I (hay fever, asthma)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: None known

Comments: This category represents at least three genera, including Bipolaris, Drechslera, and Exserohilum. This group cannot be consistently separated by spore morphology alone.

Cladosporium

Outdoor Habitat: Cladosporium is one of the most common environmental fungi observed worldwide and is widely reported from soil and decaying vegetation.

Cladosporium herbarum and C. cladosporioides are among the most frequently encountered species, both in outdoor and indoor environments.

Indoor Habitat: Wetted wood and gypsum wallboard paper, paper products, textiles, rubber, window sills. Cladosporium has the ability to grow at low temperatures and can thus, grow on rubber gaskets and food in refrigerators.

Allergy Potential: Type I (hay fever, asthma) - an important and common outdoor allergen

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals. Cladosporium are some of the most common species reported as indoor contaminants, occasionally linked to health problems.

Toxin Potential: Cladosporium has two known toxins (cladosporin and emodin). These toxins are not known to be highly toxic. There is no evidence in the literature of toxic effects associated to inhalation of Cladosporium conidia (spores) indoors.

Comments: The most commonly reported spore in the outdoor air worldwide. This makes Cladosporium one of the most commonly reported and abundant spore types both indoors and outdoors. The prevalence of this spore can vary throughout the year, but is especially high in late summer and autumn, especially where cereal crops are commonly planted.

An important and common allergen source.

Cladosporium sphaerospermum

Outdoor Habitat: Dead or dying cellulosic materials like wood and leaves.

Indoor Habitat: A favorite place for this mold type to grow is on wetted or rotting window sills. Also likes to grow on all type of wetted or rotting wood.

Allergy Potential: Type I (hay fever, asthma) - an important and common outdoor allergen

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: None known

Comments: A common mold on wetted wood, especially on window sills. Not frequently found in the air.

Curvularia

Outdoor Habitat: Soil and decaying vegetation
Indoor Habitat: Wetted wood and gypsum wallboard paper, many cellulytic substrates
Allergy Potential: Type I (hay fever, asthma), common cause of allergenic rhinitis
Disease Potential: Potential human pathogen in immunocompromised people
Toxin Potential: None known
Comments: None

Epicoccum

Outdoor Habitat: Epicoccum is a widespread cosmopolitan that grows on dead or decaying organic matter, wood, textiles, paper, a variety of foods, insects and human skin. It is commonly found in the soil. Epicoccum spores are more prevalent on dry, windy days, with higher counts late in the day.
Indoor Habitat: Capable of growing on a wide variety of substrates and manufactured products found indoors when wetted such as gypsum board, floors, carpets, mattress dust, and house plants.
Allergy Potential: Type I (hay fever, asthma)
Disease Potential: None known
Toxin Potential: None known
Comments: Very common in outdoor air in the summer months, especially in the midwest USA during harvest times.

Nigrospora

Outdoor Habitat: Soil and decaying vegetation
Indoor Habitat: Wetted wood and gypsum wallboard paper
Allergy Potential: Type I (hay fever, asthma)
Disease Potential: None known
Toxin Potential: None known
Comments: Rarely observed growing indoors

Penicillium

Outdoor Habitat: Soil and decaying vegetation, textiles, fruits

Indoor Habitat: Wetted wood and gypsum wallboard paper, textiles, leather

Allergy Potential: Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: Several known

Comments: Extremely common in indoor air, but without the fruiting bodies associated with the spores will be listed as "Penicillium / Aspergillus" group.
Penicillium identified in air samples indicates that the fruiting bodies were observed. This usually suggests that the source of the mold is nearby and / or a growth was disturbed. The fruiting bodies are not easily sent airborne, nor do they stay in the air long.

Penicillium/Aspergillus

Outdoor Habitat: Soil and decaying vegetation, textiles, fruits. These spores are commonly observed and are a normal part of outside air.

Indoor Habitat: Wetted wood and gypsum wallboard paper, textiles, leather, able to grow on many types of substrates.

Allergy Potential: Type I (hay fever, asthma), Type III (hypersensitivity pneumonitis)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: Several known

Comments: Extremely common in indoor air in low amounts. This type of spore should not constitute an overwhelming percentage and be present in very high numbers.

These two genera are grouped together because they cannot be reliably differentiated into their respective genera based solely on spore morphology.

Pithomyces

Outdoor Habitat: Soil and decaying vegetation and their spores are easily dispersed into the air by wind

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: None known

Disease Potential: None known

Toxin Potential: One known (sporidesmin)

Comments: A very common spore type in the air. Can be a water indicator mold type indoors

---

Scopulariopsis

Outdoor Habitat: Soil and decaying vegetation, dung

Indoor Habitat: Wetted wood and gypsum wallboard paper

Allergy Potential: Type III (hypersensitivity pneumonitis)

Disease Potential: Opportunistic pathogen in immunocompromised persons, not normally a pathogen in healthy individuals.

Toxin Potential: Not well studied

Comments: Easily dispersed by wind and air currents. Can grow with very little water and readily grow on wallboard when high humidity situations, e.g. closets - capable of growing on leather clothes

---

Smut/Myxomycetes

Outdoor Habitat: Soil and decaying vegetation and wood, especially dead stumps and bark

Indoor Habitat: Not known to grow indoors, sometimes found on firewood

Allergy Potential: Type I (hay fever, asthma), rare

Disease Potential: None known

Toxin Potential: None known

Comments: These two groups are difficult to distinguish due to their "round, brown" morphology. Smuts are especially common in the environment and can be seen in indoor air samples even during the winter in homes because the spores can get trapped in carpets

---